

# Regions-madogram in spatial processes

C. Fonseca <sup>\*</sup>  
Instituto Politécnico da Guarda.  
Portugal

L. Pereira, A.P. Martins, H. Ferreira <sup>†</sup>  
Departamento de Matemática  
Universidade da Beira Interior  
Portugal

---

**Abstract:** Spatial environmental processes often exhibit dependence in their large values. In order to model such processes their dependence properties must be characterized and quantified. In this paper we introduce a regions-madogram,  $\nu_F(\mathbf{A}, \mathbf{B})$ , that evaluates the dependence among extreme observations located in two separated regions,  $\mathbf{A}$  and  $\mathbf{B}$ , of  $\mathbb{Z}^2$ . We compute the range of this new dependence measure and compare it with extremal coefficients, finding generalizations of the known relations in pairwise approach. The results are illustrated in two max-stable processes: the Schlather's and the Geometric Gaussian models.

---

## 1 Introduction

Quantifying dependence is a central issue when leading with spatial observations. In conventional geostatistics, the degree of pairwise spatial dependence in observations from a max-stable random field  $\mathbf{X} = \{X_{\mathbf{t}}\}_{\mathbf{t} \in \mathbb{Z}^2}$  is determined by the variogram

$$\gamma(\mathbf{i}, \mathbf{j}) = \frac{1}{2} E (X_{\mathbf{i}} - X_{\mathbf{j}})^2.$$

If finite,  $\gamma(\mathbf{i}, \mathbf{j})$  captures some of the spatial structure, capturing it completely if the field is Gaussian. However, it is not possible to ensure that  $\gamma(\mathbf{i}, \mathbf{j})$  is finite for fields of maxima, so this second-order statistic may not be well-adapted for extremes. To ensure that the moment quantities are finite, Naveau *et al.* (2005) introduced the following type of first-order variograms,

$$\nu_F(\mathbf{i}, \mathbf{j}) = \frac{1}{2} E |F(X_{\mathbf{i}}) - F(X_{\mathbf{j}})|,$$

where  $F$  is the common distribution function of  $X_{\mathbf{i}}$ ,  $\mathbf{i} \in \mathbb{Z}^2$ .

More recently, Cooley *et al.* (2006) investigated the basic properties of first-order variograms, also called madograms, for spatial extreme fields. In particular, they derived a few relationships between the

---

<sup>\*</sup>cfonseca@ipg.pt

<sup>†</sup>lpereira@mat.ubi.pt; amartins@mat.ubi.pt; helena@mat.ubi.pt

normalized madogram,  $\nu_F$ , and the extremal coefficient defined in Schlather (2002) and Schlather and Tawn (2003) as

$$P(\max(X_{\mathbf{i}}, X_{\mathbf{j}}) \leq u) = F^{\varepsilon(\mathbf{i}, \mathbf{j})}(u).$$

The normalized madogram fully characterizes the extremal coefficient since we have

$$\varepsilon(\mathbf{i}, \mathbf{j}) = \frac{1 + 2\nu_F(\mathbf{i}, \mathbf{j})}{1 - 2\nu_F(\mathbf{i}, \mathbf{j})}, \quad (1.1)$$

or equivalently

$$\nu_F(\mathbf{i}, \mathbf{j}) = \frac{1}{2} - \frac{1}{\varepsilon(\mathbf{i}, \mathbf{j}) + 1}. \quad (1.2)$$

Despite that, in practice, the pairwise dependence measures seem to be the most useful and easily understandable, we shall introduce a regions-madogram, that evaluates the dependence among extreme observations located in two separated regions,  $\mathbf{A}$  and  $\mathbf{B}$ , of  $\mathbb{Z}^2$ .

We illustrate our results with the Geometric Gaussian and the Schlather's models for random fields.

## 2 Measuring spatial dependence for maxima over separated regions

We define a normalized madogram to assess dependence among extreme observations located in two separated regions  $\mathbf{A}$  and  $\mathbf{B}$  of  $\mathbb{Z}^2$ , as follows.

**Definition 2.1** Let  $\mathbf{X}$  be a max-stable random field and  $\mathbf{A}$  and  $\mathbf{B}$  two separated regions of  $\mathbb{Z}^2$ . The regions-madogram,  $\nu_F(\mathbf{A}, \mathbf{B})$ , is defined as

$$\nu_F(\mathbf{A}, \mathbf{B}) = \frac{1}{2|\mathbf{A}||\mathbf{B}|} \sum_{\mathbf{i} \in \mathbf{A}} \sum_{\mathbf{j} \in \mathbf{B}} E|F(X_{\mathbf{i}}) - F(X_{\mathbf{j}})|,$$

or equivalently

$$\nu_F(\mathbf{A}, \mathbf{B}) = \frac{1}{|\mathbf{A}||\mathbf{B}|} \sum_{\mathbf{i} \in \mathbf{A}} \sum_{\mathbf{j} \in \mathbf{B}} \nu_F(\mathbf{i}, \mathbf{j})$$

**Remark 2.1** To measure the dependence among the variables  $X_{\mathbf{i}}$  and  $X_{\mathbf{j}}$ ,  $\mathbf{j} \in V^{(n)}(\mathbf{i})$ , where

$$V^{(n)}(\mathbf{i}) = \{\mathbf{j} \in \mathbb{Z}^2 : \max\{|i_s - j_s| : s = 1, 2\} = n\},$$

we can consider the sequence

$$h(n) = \nu_F(\mathbf{i}, n) = \nu_F(\mathbf{i}, V^{(n)}(\mathbf{i})).$$

**Remark 2.2** The regions-madogram,  $\nu_F(\mathbf{A}, \mathbf{B})$ , take values in  $[0, \frac{1}{6}]$ .

From the definition of regions-madogram we can establish the following properties.

**Proposition 2.1** *Suppose that  $\mathbf{X}$  is a max-stable random field.*

1. Let  $\mathbf{A} \subseteq \mathbb{Z}^2$  and  $\mathbf{A} + \mathbf{s} = \{\mathbf{i} + \mathbf{s} : \mathbf{i} \in \mathbf{A}\}$ ,  $\mathbf{s} \in \mathbb{Z}^2$ . If  $\mathbf{X}$  is stationary, then

$$\nu_F(\mathbf{A} + \mathbf{s}, \mathbf{B} + \mathbf{s}) = \nu_F(\mathbf{A}, \mathbf{B})$$

2. Let  $\mathbf{A}_i$ ,  $i = 1, \dots, p$ , be disjoint subsets of  $\mathbb{Z}^2$ . Then

$$\nu_F\left(\bigcup_{i=1}^p \mathbf{A}_i, \mathbf{B}\right) = \sum_{i=1}^p \alpha_i \nu_F(\mathbf{A}_i, \mathbf{B}),$$

with  $\alpha_i = \frac{|\mathbf{A}_i|}{|\bigcup_{i=1}^p \mathbf{A}_i|}$ .

In particular,

$$\nu_F(\mathbf{A}, \mathbf{B}) = \sum_{\mathbf{i} \in \mathbf{A}} \frac{1}{|\mathbf{A}|} \nu_F(\{\mathbf{i}\}, \mathbf{B}).$$

The normalized regions-madogram is related to the pairwise extremal coefficients  $\varepsilon(\mathbf{i}, \mathbf{j})$ ,  $\mathbf{i} \in \mathbf{A}$ ,  $\mathbf{j} \in \mathbf{B}$ , through

$$\nu_F(\mathbf{A}, \mathbf{B}) = \frac{1}{2} - \frac{1}{|\mathbf{A}| |\mathbf{B}|} \sum_{\mathbf{i} \in \mathbf{A}} \sum_{\mathbf{j} \in \mathbf{B}} \frac{1}{\varepsilon(\mathbf{i}, \mathbf{j}) + 1}.$$

In the following, we define a new dependence measure,  $\varepsilon^*(\mathbf{A}, \mathbf{B})$ , that preserves relation (1.1).

**Definition 2.2** *Let  $\mathbf{X}$  be a max-stable random field and  $\mathbf{A}, \mathbf{B} \subseteq \mathbb{Z}^2$ . The coefficient  $\varepsilon^*(\mathbf{A}, \mathbf{B})$  is defined as*

$$\varepsilon^*(\mathbf{A}, \mathbf{B}) = \frac{1}{\frac{1}{|\mathbf{A}| |\mathbf{B}|} \sum_{\mathbf{i} \in \mathbf{A}} \sum_{\mathbf{j} \in \mathbf{B}} \frac{1}{\varepsilon(\mathbf{i}, \mathbf{j}) + 1}} - 1. \quad (2.1)$$

We have  $\varepsilon^*(\mathbf{A}, \mathbf{B}) = \frac{1+2\nu_F(\mathbf{A}, \mathbf{B})}{1-2\nu_F(\mathbf{A}, \mathbf{B})}$ .

**Remark 2.3** The coefficient  $\varepsilon^*(\mathbf{A}, \mathbf{B})$  take values in  $[1, 2]$ . When  $\varepsilon^*(\mathbf{A}, \mathbf{B})=1$ , we have the complete dependence for each pair of variables  $(X_{\mathbf{i}}, X_{\mathbf{j}})$ ,  $\mathbf{i} \in \mathbf{A}, \mathbf{j} \in \mathbf{B}$ , and conversely. For each  $\mathbf{i} \in \mathbf{A}$  and  $\mathbf{j} \in \mathbf{B}$ , the variables  $X_{\mathbf{i}}$  and  $X_{\mathbf{j}}$  are independent if and only if  $\varepsilon^*(\mathbf{A}, \mathbf{B}) = 2$ .

### 3 Applications

In this section we illustrate our results in well known max-stable processes: the Schlather's model and the Geometric Gaussian model.

**Example 3.1 (Schlather's model)** In Schlather (2002), a new class of bivariate marginal distributions is defined, by the extremal Gaussian process, in the following way

$$P(X_{\mathbf{i}} \leq x, X_{\mathbf{j}} \leq y) = \exp \left[ -\frac{1}{2} \left( \frac{1}{x} + \frac{1}{y} \right) \left( 1 + \sqrt{1 - \frac{2(\rho(\|\mathbf{i} - \mathbf{j}\|) + 1)xy}{(x+y)^2}} \right) \right], \quad x, y \in \mathbb{R},$$

where  $\rho(\cdot)$  is the covariance function of the underlying Gaussian process.

We will consider that  $\rho(\|\mathbf{i} - \mathbf{j}\|) = \exp(-(\|\mathbf{i} - \mathbf{j}\|))$ .




Figure 3.1. Schlather's model with  $\rho(\|\mathbf{i} - \mathbf{j}\|) = \exp(-(\|\mathbf{i} - \mathbf{j}\|))$

The extremal coefficient is given by

$$\varepsilon(\mathbf{i}, \mathbf{j}) = 1 + \sqrt{\frac{1 - \rho(\|\mathbf{i} - \mathbf{j}\|)}{2}}$$

and the amount of dependence among  $X_{\mathbf{i}}$  and the variables  $X_{\mathbf{j}}, \mathbf{j} \in V^{(n)}(\mathbf{i})$ , is given by

$$\begin{aligned} h(n) &= \nu_F(\mathbf{i}, V^{(n)}(\mathbf{i})) \\ &= \frac{1}{2} - \frac{1}{8n} \left( \frac{4}{2 + \sqrt{\frac{1 - \exp(-n\sqrt{2})}{2}}} + \frac{4}{2 + \sqrt{\frac{1 - \exp(-n)}{2}}} + \sum_{i=1}^{n-1} \frac{8}{2 + \sqrt{\frac{1 - \exp(-\sqrt{n^2+i^2})}{2}}} \right). \end{aligned}$$




Figure 3.2. Quantifying dependence between the regions  $\{\mathbf{i}\}$  and  $V^{(n)}(\mathbf{i})$  in the Schlather's model

**Example 3.2 (Geometric Gaussian model)** Let  $\mathbf{X}$  be a Geometric Gaussian random field. The bivariate distribution is given by

$$P(X_{\mathbf{i}} \leq x, X_{\mathbf{j}} \leq y) = \exp \left( -\frac{1}{x} \Phi \left( \frac{a}{2} + \frac{1}{a} \log \frac{y}{x} \right) - \frac{1}{y} \Phi \left( \frac{a}{2} + \frac{1}{a} \log \frac{x}{y} \right) \right), \quad x, y \in \mathbb{R},$$

where  $\Phi$  is the standard normal distribution function and

$$a^2 = 2\sigma^2(1 - \rho(\|\mathbf{i} - \mathbf{j}\|)),$$

where  $\rho(\cdot)$  is the covariance function of the Gaussian process. Let us consider  $\sigma = 1$ ,  $\mathbf{A} = \{\mathbf{i}, \mathbf{j}\}$  and  $\mathbf{B} = \{\mathbf{i} + (s, 0), \mathbf{j} + (s, 0)\}$ ,  $s \in \mathbb{N}$ .

We have

$$\varepsilon(\mathbf{i}, \mathbf{j}) = 2\Phi \left( \sqrt{\frac{1 - \rho(\sqrt{(i_1 - j_1)^2 + (i_2 - j_2)^2})}{2}} \right)$$

and

$$\begin{aligned} \nu_F(\mathbf{A}, \mathbf{B}) = & \frac{1}{2} - \frac{1}{4} \left( \frac{1}{1 + \Phi \left( \sqrt{\frac{1 - \rho(|s|)}{2}} \right)} + \frac{1}{1 + 2\Phi \left( \sqrt{\frac{1 - \rho(\sqrt{(-j_1 + i_1 + s)^2 + (i_2 - j_2)^2})}{2}} \right)} \right. \\ & \left. + \frac{1}{1 + 2\Phi \left( \sqrt{\frac{1 - \rho(\sqrt{(j_1 - i_1 + s)^2 + (i_2 - j_2)^2})}{2}} \right)} \right). \end{aligned}$$

## References

- [1] Cooley, D., Naveau, P., and Poncet, P. (2006). Variograms for spatial max-stable random fields. In Springer, editor, *Dependence in Probability and Statistics*, Volume 187, 373–390. Springer, New York, Lecture Notes in Statistics Edition.
- [2] Naveau, P., Cooley, D., and Pocet, P. (2005). Variograms for spatial max-stable random fields. *Manuscript*.
- [3] Ribatet, M. (2009). A users guide to the spatialextremes package. Technical report, École Polytechnique Fédérale de Lausanne.
- [4] Schlather, M. (2002). Models for stationary max-stable random fields. *Extremes*, 5(1), 3344.
- [5] Schlather, M. and Tawn, J. (2003). A dependence measure for multivariate and spatial extremes: Properties and inference. *Biometrika*, 90(1), 139-156.
- [6] Smith, R. (1990). Max-stable processes and spatial extremes. Unpublished manuscript.